

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/poamed>

Original research article

Lack of association between two genetic polymorphisms of SOD2 (rs2758339 and rs5746136) and the risk of opium dependency



Fariba Boroumand, Mostafa Saadat*

Department of Biology, College of Sciences, Shiraz University, Shiraz 71467-13565, Iran

ARTICLE INFO

Article history:

Received 21 January 2017

Received in revised form

30 January 2017

Accepted 23 March 2017

Available online 6 July 2017

Keywords:

SOD2

Opium dependency

rs2758339

rs5746136

ABSTRACT

Introduction: Superoxide dismutase-2 (EC 1.15.1.1; SOD2, OMIM: 147460) is a tetrameric enzyme which contains manganese in its active site. It is an important enzyme involved in the cellular detoxification by converting highly toxic superoxide radicals into less reactive molecules, hydrogen peroxide and oxygen. Several single nucleotide polymorphisms have been well defined in the gene encoding SOD2, including the potentially functional polymorphisms of rs2758339 and rs5746136.

Aim: The aim of the present study is to investigate the associations between the rs2758339 (A/C substitution) and rs5746136 (A/G substitution) polymorphisms and the risk of dependency to opium.

Material and methods: The present case-control study was performed in Shiraz (southern Iran) on 143 opium dependent and 569 healthy controls. The genotypes of the rs2758339 and rs5746136 polymorphisms were determined by polymerase chain reaction.

Results and discussion: Statistical analysis showed that there was no significant association between the study polymorphisms and the risk of opium dependency. A significant linkage disequilibrium was observed between the study SOD2 polymorphisms. Statistical analysis showed that there was no significant association between the haplotypes of the polymorphisms and the risk of opium dependency.

Conclusions: The present data revealed that the rs2758339 and rs5746136 polymorphisms of SOD2 are not risk factors for dependency to opium.

© 2017 Published by Elsevier Sp. z o.o. on behalf of Warmińsko-Mazurska Izba Lekarska w Olsztynie.

1. Introduction

Superoxide dismutase-2 (EC 1.15.1.1; SOD2, OMIM: 147460) is a tetrameric enzyme which contains manganese in its active

site.^{1,2} SOD2 metabolizes superoxide radicals into molecular oxygen and hydrogen peroxide (H₂O₂)³; subsequently H₂O₂ is converted into water by other enzymes such as catalase. SOD2 is expressed in most tissues, including in human brain.⁴ In human, several single genetic polymorphisms have been

* Corresponding author. Department of Biology, College of Sciences, Shiraz University, Shiraz, 71454 Iran. Tel.: +9871 36137432; fax: +9871 32280926.

E-mail addresses: fariba.boroumand@gmail.com (F. Boroumand), saadat@shirazu.ac.ir, msaadat41@yahoo.com (M. Saadat).

reported in the *SOD2* (https://www.ncbi.nlm.nih.gov/projects/SNP/snp_ref.cgi?geneid=6648), including the rs2758339 (A/C substitution)⁵ and rs5746136 (A/G substitution) polymorphisms.⁴ These polymorphisms are located in the vicinity of SP1 and NF- κ B transcription element sequences⁴ and glucocorticoid receptor binding site.⁵ Association between these *SOD2* polymorphisms and the risks of several multifactorial complex traits has been studied by researchers.^{5–12}

It has been well demonstrated that opiates may cause oxidative stress in drug dependent persons.^{13–15} It has been reported that morphine decreases enzyme activities of the superoxide dismutase and catalase.^{16–18} On the other hand the mRNA levels of the *SOD2* decreased when SH-SH5Y cells were treated by methadone and morphine for a long period.^{19,20} These reports indicated that drug-abused persons experience the oxidative stress.

The association between some polymorphisms of the genes involved in cellular detoxification (such as members of glutathione S-transferases superfamily, catalase) and susceptibility to methamphetamine and heroin abuse has been reported.^{21–26} However, there are no study investigating the association between the *SOD2* polymorphisms and risk of opium dependency. These facts sufficiently provide us with a theoretical ration to carry out the present case–control study.

2. Aim

This study was conducted to determine the association of the rs2758339 (A/C substitution) and rs5746136 (A/G substitution) genetic polymorphism of *SOD2* with the risk of opium dependency in Iranian population.

3. Material and methods

3.1. Ethical approval

This study was approved by the Shiraz University ethics committee. Informed consent was obtained from each subject before the study. This work was carried out in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki) for Ethical Principles for Medical Research Involving Human Subjects.

3.2. Subjects

A detailed description of the subjects has been reported in our previous article.²⁴ In brief, this case–control study was conducted in Shiraz (southern Iran) on 143 (12 females, 131 males) opium dependent persons and 570 (55 females, 515 males) healthy controls. The patients were in methadone maintenance for treating their dependency and all of them reported opium as their primary drug of choice. Control individuals were blood donors, who declared that they did not suffer from substance abuse. The mean age (SD) of the patients and the controls were 39.6 (11.3) and 40.8 (10.7) years, respectively. There was no significant difference between the two study groups for their gender and age distribution ($P > 0.05$). Iranian populations showed high level of heterogeneity,^{27–29}

therefore, we selected the participants from Persian Muslims (Caucasians) living in Shiraz (Fars province, southern Iran).

3.3. Genotyping

Genomic DNA was isolated from EDTA treated blood samples. The polymerase chain reaction (PCR) conditions for determining the genotypes of the polymorphisms and quality control were the same as that reported previously.^{3,4} It should be noted that we failed to successfully determine the polymorphisms in two participants of control group, explaining the variation in the total number of samples listed in Table 1.

3.4. Statistical analysis

Goodness-of-fit χ^2 test was used to verify whether the observed genotypic frequencies were in accordance with the Hardy–Weinberg equilibrium. The associations between the genotypes and the risk of opium dependency were assessed by odds ratios (ORs) and 95% confidence intervals (CIs). The software SNPalyze(TM) ver. 6 Standard (Dynacom Co, Ltd. Kanagawa, Japan) was used to evaluate the status of pair wise linkage disequilibrium for the studied polymorphisms. Statistical analysis was performed using SPSS software (Chicago, IL, USA; version 11.5). A probability of $P < 0.05$ was considered statistically significant.

4. Results

The genotypic frequency of the study polymorphisms in the opium dependent cases and healthy controls is shown in Table 1. The genotypic frequencies of the study polymorphisms in healthy controls (for rs2758339 polymorphism: $\chi^2 = 0.32$, $df = 1$, $P = 0.566$; for rs5746136 polymorphism: $\chi^2 = 0.01$, $df = 1$, $P = 0.928$) were consistent with the Hardy–Weinberg equilibrium distribution. Statistical analysis showed that there was no significant association between the study polymorphisms and the risk of opium dependency (Table 1). There was no liner trend for the numbers of putative high risk alleles of the polymorphisms and the risk of opium dependency (for rs2758339 polymorphism: $\chi^2 = 0.33$, $P = 0.565$; for rs5746136 polymorphism: $\chi^2 = 0.43$, $P = 0.509$).

A significant linkage disequilibrium was observed between the study polymorphisms (for control group: $D' = -0.9375$, $r^2 = 0.3994$, $\chi^2 = 455.2$, $P < 0.001$; for opium dependent group: $D' = -1.0$, $r^2 = 0.4486$, $\chi^2 = 128.2$, $P < 0.001$). Table 2 shows the haplotypic frequencies of the study polymorphisms in opium dependent patient and control groups. The CG haplotype was used as a reference. Statistical analysis showed that there was no significant association between the haplotypes of rs2758339 and rs5746136 polymorphisms and the risk of opium dependency (Table 2).

5. Discussion

Based on the findings of several lines of evidences, including serum markers of inflammation¹³ and frequency of micro-nuclei in drug abused persons,¹⁵ and also alterations of

Table 1 – Association between the rs2758339 and rs5746136 genetic polymorphisms of SOD2 and the risk of dependency to opium.

Genotypes	Controls (%)	Cases (%)	OR	95% CI	P value
rs2758339 polymorphism					
AA	180 (31.6)	43 (30.1)	1.0	–	–
AC	286 (50.3)	71 (49.7)	1.03	0.68–1.58	0.858
CC	103 (18.1)	29 (20.2)	1.17	0.69–2.00	0.543
Dominant model					
AA	180 (31.6)	43 (30.1)	1.0	–	–
AC + CC	389 (68.4)	100 (69.9)	1.07	0.72–1.60	0.718
Recessive model					
AA + AC	466 (81.9)	114 (79.7)	1.0	–	–
CC	103 (18.1)	29 (20.3)	1.15	0.72–1.82	0.549
Alleles					
A	646 (56.8)	157 (54.9)	1.0	–	–
C	492 (43.2)	129 (45.1)	1.07	0.83–1.40	0.568
rs5746136 polymorphism					
GG	222 (39.1)	58 (40.5)	1.0	–	–
GA	267 (47.0)	69 (48.3)	0.98	0.66–1.46	0.956
AA	79 (13.9)	16 (11.2)	0.77	0.42–1.42	0.413
Dominant model					
GG	222 (39.1)	58 (40.5)	1.0	–	–
GA + AA	346 (60.9)	85 (59.5)	0.94	0.64–1.36	0.747
Recessive model					
GG + GA	489 (86.1)	127 (88.8)	1.0	–	–
AA	79 (13.9)	16 (11.2)	0.78	0.44–1.38	0.394
Alleles					
G	711 (62.6)	185 (64.7)	1.0	–	–
A	425 (37.4)	101 (35.3)	0.91	0.69–1.19	0.511

OR: odds ratio; CI: confidence interval.

Table 2 – Association between haplotypes of the rs2758339 and rs5746136 genetic polymorphisms of SOD2 and the risk of dependency to opium.

Haplotypes		Controls	Cases	OR	95% CI	P value
rs2758339	rs5746136					
C	G	479	129	1.0	–	–
A	A	413	101	0.90	0.67–1.21	0.517
A	G	232	56	0.89	0.63–1.27	0.541
C	A	12	0	–	–	–

OR: odds ratio; CI: confidence interval.

enzyme activities of SOD and catalase^{16–18} or alterations of the SOD2 mRNA levels in experimental animals or cells after exposure by morphine or methadone,^{19,20} it might be concluded that drug dependent persons are under oxidative stress. On the other hand, oxidative stress is very important in brain, because it has effects not only on the integrity of DNA but also has some effect(s) on the function of the N-type and/or L-Type Ca²⁺ channel.³⁰ SOD2 is an important enzyme involved in the cellular detoxification. It can convert highly toxic superoxide radicals into less reactive molecules, hydrogen peroxide and oxygen.^{1,3} There are evidences that several polymorphisms are associated with alteration in the mRNA levels of the corresponding genes.^{31–33} Taken together we hypothesized that genetic polymorphisms in SOD2, may be associated with the risk of dependency to opium, by its influence on the SOD2 expression levels. However, the present study did not support our hypothesis. The study polymorphisms are located in the vicinity of SP1 and NF- κ B transcription element sequences⁴ and glucocorticoid receptor binding

site.⁵ Therefore, it is necessary to investigate the influence of these polymorphisms on the SOD2 mRNA levels in future studies.

There are some limitations of this study. First, the number of opium dependent persons was relatively small. Second, the class of superoxide dismutase contain 3 members (SOD1, SOD2, and SOD3). In the present study we study polymorphisms of the SOD2. Third although it has been reported that the SOD2 is very polymorphic, here we study only two of them. Considering that the association between polymorphisms and a particular trait may show significant different patterns between ethnic groups,^{34–36} the other studies should carry out in other populations.

6. Conclusion

It should be noted that it is the first association study between the rs2758339 and rs5746136 polymorphisms and the risk of

opium dependency. The present data revealed that the rs2758339 and rs5746136 polymorphisms of SOD2 are not risk factors for dependency to opium.

Conflict of interest

None declared.

Contributors

The first author was involved in data collection and statistical analyses the second author was involved in designed the study, statistical analyses. Both authors wrote the manuscript and have approved the final manuscript.

Acknowledgments

The authors are indebted to the participants for their close cooperation. The authors are indebted to Dr. Maryam Ansari-Lari for her critical reading the manuscript and for her contribution in discussion. This study was supported by Shiraz University, Iran.

REFERENCES

- Beckman G, Lundgren E, Tarnvik A. Superoxide dismutase isozymes in different human tissues, their genetic control and intracellular localization. *Hum Hered.* 1973;23:338–345.
- Beck Y, Oren R, Amit B, et al. Human Mn superoxide dismutase cDNA sequence. *Nucleic Acids Res.* 1987;15:9076.
- Niwa J, Yamada S, Ishigaki S, et al. Disulfide bond mediates aggregation, toxicity, and ubiquitylation of familial amyotrophic lateral sclerosis-linked mutant SOD1. *J Biol Chem.* 2007;282:28087–28095.
- Wan XS, Devalaraja MN, St Clair DK. Molecular structure and organization of the human manganese superoxide dismutase gene. *DNA Cell Biol.* 1994;13:1127–1136.
- Hernandez-Saavedra D, McCord JM. Association of a new intronic polymorphism of the SOD2 gene (G1677T) with cancer. *Cell Biochem Funct.* 2009;27:223–227.
- Ebrahimpour S, Saadat I. Association of CAT C-262T and SOD1 A251G single nucleotide polymorphisms susceptible to gastric cancer. *Mol Biol Res Commun.* 2014;3:223–229.
- Zhou Y, Shuai P, Li X, et al. Association of SOD2 polymorphisms with primary open angle glaucoma in a Chinese population. *Ophthalmic Genet.* 2015;36:43–49.
- Poggi C, Giusti B, Vestri A, et al. Genetic polymorphisms of antioxidant enzymes in preterm infants. *J Matern Fetal Neonatal Med.* 2012;25(suppl 4):131–134.
- Ding G, Liu F, Shen B, et al. The association between polymorphisms in prooxidant or antioxidant enzymes (myeloperoxidase, SOD2, and CAT) and genes and prostate cancer risk in the Chinese population of Han nationality. *Clin Genitourin Cancer.* 2012;10:251–255.
- Giusti B, Vestri A, Poggi C, et al. Genetic polymorphisms of antioxidant enzymes as risk factors for oxidative stress-associated complications in preterm infants. *Free Radic Res.* 2012;46:1130–1139.
- Park HY, Kim JH, Lim YH, et al. Influence of genetic polymorphisms on the association between phthalate exposure and pulmonary function in the elderly. *Environ Res.* 2013;122:18–24.
- Wiener HW, Perry RT, Chen Z, et al. A polymorphism in SOD2 is associated with development of Alzheimer's disease. *Genes Brain Behav.* 2007;6:770–775.
- Cunha-Oliveira T, Rego AC, Oliveira CR. Cellular and molecular mechanisms involved in the neurotoxicity of opioid and psychostimulant drugs. *Brain Res Rev.* 2008;58:192–208.
- Ghazavi A, Mosayebi G, Solhi H, et al. Serum markers of inflammation and oxidative stress in chronic opium (Taryak) smokers. *Immunol Lett.* 2013;153:22–26.
- Soykut B, Eken A, Erdem O, et al. Oxidative stress enzyme status and frequency of micronuclei in heroin addicts in Turkey. *Toxicol Mech Methods.* 2013;23:684–688.
- Payabvash S, Beheshtian A, Salmasi AH, et al. Chronic morphine treatment induces oxidant and apoptotic damage in the mice liver. *Life Sci.* 2006;79:972–980.
- Mao J, Sung B, Ji R, et al. Neuronal apoptosis associated with morphine tolerance: evidence for an opioid-induced neurotoxic mechanism. *J Neurosci.* 2002;22:7650–7661.
- Zhou J, Li Y, Yan G, et al. Protective role of taurine against morphine-induced neurotoxicity in C6 cells via inhibition of oxidative stress. *Neurotox Res.* 2011;4:334–342.
- Saify K, Saadat I, Saadat M. Down-regulation of antioxidant genes in human SH-SY5Y cells after treatment with morphine. *Life Sci.* 2016;144:26–29.
- Saify K, Saadat M. Expression patterns of antioxidant genes in human SH-SY5Y cells after treatment with methadone. *Psychiatry Res.* 2015;230:116–119.
- Nakatome M, Miyaji A, Mochizuki K, et al. Association between the GST genetic polymorphisms and methamphetamine abusers in the Japanese population. *Leg Med (Tokyo).* 2009;11(suppl 1):S468–S470.
- Koizumi H, Hashimoto K, Kumakiri C, et al. Association between the glutathione S-transferase M1 gene deletion and female methamphetamine abusers. *Am J Med Genet B: Neuropsychiatr Genet.* 2004;126B:43–45.
- Khalighinasab MR, Saify K, Saadat M. Association between GSTM1 and GSTT1 polymorphisms and susceptibility to methamphetamine dependence. *Mol Biol Res Commun.* 2015;4:25–32.
- Khalighinasab MR, Saify K, Saadat M. Association between null alleles of GSTM1 and GSTT1 and dependence to heroin and opium. *Psychiatry Res.* 2015;228:977–978.
- Hashimoto T, Hashimoto K, Miyatake R, et al. Association study between polymorphisms in glutathione-related genes and methamphetamine use disorder in a Japanese population. *Am J Med Genet B: Neuropsychiatr Genet.* 2008;147B:1040–1046.
- Saify K, Khalighinasab MR, Saadat M. No association between GSTM1 and GSTT1 genetic polymorphisms and susceptibility to opium sap dependence. *Mol Biol Res Commun.* 2016;5:59–64.
- Rafiee L, Saadat I, Saadat M. Glutathione S-transferase genetic polymorphisms (GSTM1, GSTT1 and GSTO2) in three Iranian populations. *Mol Biol Rep.* 2010;37:155–158.
- Saadat M. Distribution of ACE insertion/deletion (I/D) polymorphism in Iranian populations. *Mol Biol Res Commun.* 2015;4:63–66.
- Fallahzadeh-Abarghoeei L, Zahedi T, Mirabedi F, et al. Allelic prevalence of intron 3 insertion/deletion genetic polymorphism of DNA double-strand break repair gene XRCC4 in four Iranian populations. *Egypt J Med Hum Genet.* 2015;16:215–218.
- Hool LC. Evidence for the regulation of L-type Ca²⁺ channels in the heart by reactive oxygen species: mechanism for

- mediating pathology. *Clin Exp Pharmacol Physiol.* 2008;35:229-234.
31. Saify K, Saadat I, Saadat M. Influence of A-21T and C-262T genetic polymorphisms at the promoter region of the catalase (CAT) on gene expression. *Environ Health Prev Med.* 2016;21:382-386.
 32. Saify K, Khalighinasab MR, Saadat M. Influence of a 50 bp Ins/Del polymorphism at promoter of the superoxide dismutase-1 on gene expression and risk of heroin dependency. *Environ Health Prev Med.* 2017;22:4.
 33. Rajaei M, Saadat I, Saadat M. The novel allele (3R) of the VNTR polymorphism in the XRCC5 promoter region dramatically decreases the gene expression. *Biochem Biophys Res Commun.* 2013;430:640-641.
 34. Kiyohara C, Takayama K, Nakanishi Y. Association of genetic polymorphisms in the base excision repair pathway with lung cancer risk: a meta-analysis. *Lung Cancer.* 2006;54:267-283.
 35. Saadat M, Ansari-Lari M. Polymorphism of XRCC1 (at codon 399) and susceptibility to breast cancer, a meta-analysis of the literatures. *Breast Cancer Res Treat.* 2009;115:137-144.
 36. Saadat M. Apolipoprotein E (APOE) polymorphisms and susceptibility to breast cancer: a meta-analysis. *Cancer Res Treat.* 2012;44:121-126.